

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

### IN THE CLAIMS

Please amend the claims of the present application under the provisions of 37 C.F.R. §1.121(c), as indicated below:

1. (Currently amended): A modified release dosage form comprising of a high solubility active ingredient prepared by using dual retard technique to control the release of the high solubility active ingredient, said dosage form comprising a) micro matrix particles containing active ingredient(s) and one or more hydrophobic release ~~controlling~~ controlling agent and b) a coating of one or more hydrophobic release ~~controlling~~ controlling agents on said micro matrix particles.
2. (Original): A dosage form as claimed in claim 1, is in the form of tablet.
3. (Currently Amended): A dosage form according to claim 1, wherein the hydrophobic release controlling agents are selected from the group ~~comprising~~ consisting of ammonio methacrylate copolymers type A and B as described in USP, methacrylic acid copolymer type A, B and C as described in USP, polyacrylate dispersion 30% as described in Ph. Eur., polyvinyl acetate dispersion, ethylcellulose, cellulose acetate, cellulose propionate (lower, medium or higher molecular weight), cellulose acetate propionate, cellulose acetate butyrate, cellulose acetate phthalate, cellulose triacetate, poly(methyl methacrylate), poly(ethyl methacrylate), poly(butyl methacrylate), poly(isobutyl methacrylate), poly(hexyl methacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl actylate), poly(octadecyl acrylate), waxes ~~such as~~ selected from the group consisting of beeswax, carnauba wax, microcrystalline wax, and ozokerite; fatty alcohols ~~such as~~ selected from the group consisting of cetostearyl alcohol, stearyl alcohol; cetyl

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

alcohol and myristyl alcohol; and fatty acid esters ~~such as~~ selected from the group consisting of glyceryl monostearate, glycerol monooleate, acetylated monoglycerides, tristearin, tripalmitin, cetyl esters wax, glyceryl palmitostearate, glyceryl behenate and hydrogenated castor oil.

4. (Currently amended): A dosage form according to claim 3, wherein said hydrophobic release controlling agents are selected ~~preferably~~ from ammonio methacrylate co-polymers.

5. (Currently amended): A dosage form according to claim 4, wherein ~~preferred~~ ammonio methacrylate co-polymers are selected from the group consisting of Eudragit Ammonio Methacrylate Copolymer type B USP, Eudragit RL ~~[[()]]~~ Ammonio Methacrylate Copolymer type A USP~~[[()]]~~ and ~~Eudragit NE30D~~ ~~[[()]]~~ Polyacrylate dispersion 30% Ph. Eur.~~[[()]]~~

6. (Original): A dosage form according to claim 1, wherein in micro matrix particles, the active ingredient and one or more hydrophobic release controlling agents are present in a ratio of from 100:1 to 100:75.

7. (Currently amended): A dosage form according to claim 6, wherein in micro matrix particles, the active ingredient and one or more hydrophobic release controlling agents are present ~~preferably~~ in a ratio of from 100:2.5 to 100:50.

8. (Currently amended): A dosage form according to claim 6, wherein in micro matrix particles, the active ingredient and one or more hydrophobic release controlling agents are present ~~more preferably~~ in a ratio of from 100:2.5 to 100:30.

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

9. (Currently amended): A dosage form according to claim 6, wherein in micro matrix particles, the active ingredient and one or more hydrophobic release controlling agents are present ~~most preferably~~ in a ratio of from 100:2.5 to 100:20.

10. (Currently amended): A dosage form according to ~~claim one of claims 6-9~~ claim 6 to 9, wherein in micro matrix particles, the active ingredient can be less than or equal to 1500 mg.

11. (Currently amended): A dosage form according to claim 1, wherein a coating of one or more hydrophobic release controlling agents on said micro matrix particles ~~the hydrophobic release controlling agents employed for coating the said micro matrix particles~~ are selected from the group comprising of ammonio methacrylate copolymers type A and B as described in USP, methacrylic acid copolymer type A, B and C as described in USP, polyacrylate dispersion 30% ~~as described in Ph. Eur.~~, polyvinyl acetate dispersion, ethylcellulose, cellulose acetate, cellulose propionate (lower, medium or higher molecular weight), cellulose acetate propionate, cellulose acetate butyrate, cellulose acetate phthalate, cellulose triacetate, poly(methyl methacrylate), poly(ethyl methacrylate), poly(butyl methacrylate), poly(isobutyl methacrylate), poly(hexyl methacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate), waxes such as selected from the group consisting of beeswax, carnauba wax, microcrystalline wax, and ozokerite; fatty alcohols such as selected from the group consisting of cetostearyl alcohol, stearyl alcohol; cetyl alcohol and myristyl alcohol; and fatty acid esters such as selected from the group consisting of glyceryl monostearate; glycerol monooleate, acetylated monoglycerides, tristearin, tripalmitin, cetyl esters wax, glyceryl palmitostearate, glyceryl behenate and hydrogenated castor oil.

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

12. (Currently amended): A dosage form according to claim 11, wherein the hydrophobic release controlling agents ~~[[is]]~~ are selected from fatty acid esters.

13. (Currently amended): A dosage form according to claim 12, wherein ~~the hydrophobic release controlling agents~~ the fatty ester is are selected from the group ~~comprising~~ consisting of hydrogenated castor oil and glycerol distearate.

14. (Currently amended): A dosage form according to claim 1, wherein micro matrix particles and coating of one or more hydrophobic release controlling agents are present in a ratio of from 100:0.5 to 100:75.

15. (Currently amended): A dosage form according to claim 14, wherein micro matrix particles and coating of one or more hydrophobic release controlling agents are preferably present in a ratio of from 100:1 to 100:50.

16. (Currently amended): A dosage form according to claim 14, wherein micro matrix particles and coating of one or more hydrophobic release controlling agents are ~~more~~ preferably present in a ratio of from 100:2.5 to 100:20.

17. (Currently amended): A dosage form according to claim 1, wherein the high solubility active ingredient is selected from the group ~~comprising~~ consisting of antidiabetic agents, anti-histamines, anti-depressants, anti-viral agents, anesthetics, antacids, anti-arthritics, antibiotics, anti-psychotics, anti-spasmodics, anxiolytic agents, appetite suppressants, cardiovascular agents, cough suppressants, emollients, gastro-intestinal agents, growth regulators, respiratory stimulants, vitamins, angiotensin converting enzyme inhibitors, anti-asthmatics, anti-cholesterolemics, anti-convulsants, anti-depressants, anti-diarrhea preparations, anti-infective, anti-inflammatory agents, anti-nauseants, anti-stroke agents,

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

anti-tumor drugs, anti-tussives, anti-uricemic drugs, amino-acid preparations, antiemetics, antiobesity drugs, antiparasitics, antipyretics, appetite stimulants, cerebral dilators, chelating agents, cholecystokinin antagonists, cognition activators, deodorants, dermatological agents, diuretics, erythropoietic drugs, fertility agents, synthetic hormones, laxatives, mineral supplements, neuroleptics, neuromuscular agents, peripheral vaso-dilators, prostaglandins, vaginal preparations, vaso-constrictors, vertigo agents, biguanides, sulphonylurease, meglitinides, PPAR gama agonist [insulin sensitisers (thiazolidinedione)] and alpha-glucosidase inhibitors and the like.

18. (Original): A dosage form according to claim 1, wherein the high solubility active ingredient is selected from the group comprising of metformin hydrochloride, phenformin, buformin, captopril, ranitidine hydrochloride, potassium chloride, clindamycin, hydroxyurea, erythromycin lactobionate, vancomycin hydrochloride, balsalazide disodium, aminocaproic acid, lisinopril, tramadol, acetaminophen, ciprofloxacin, esters of ampicillin, sodium valproate, niacin, diltiazem, venlafaxine, isosorbide 5-imononitrate, isosorbide dinitrate, pentoxifylline, propranolol and quetiapine or pharmaceutically acceptable salts thereof.

19. (Original): A dosage form according to claim 1, wherein the dissolution of high solubility active ingredient is not more than 50% in 1 hour and from 25 to 90% is released in six hours.

20. (Original): A dosage form according to claim 1, wherein it reduces the chances of dose dumping, unnecessary burst effects and failure of the system.

21. (Currently amended): A dosage form according to one of claims 1 [[to]] - 18, wherein the high solubility active ingredient is potent.

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

22. (Original): A dosage form according to claim 21, wherein it reduces the chances of dose dumping, unnecessary burst effects and failure of the system.

23. (Currently amended): A dosage form according to claim 21, where the high solubility potent active ingredient is selected from the group ~~comprising~~ consisting of antidiabetic agents, antihistamines, anti-depressants, anti-viral agents, anesthetics, antacids, anti-arthritics, antibiotics, anti-psychotics, anti-spasmodics, anxiolytic agents, appetite suppressants, cardiovascular agents, cough suppressants, emollients, gastro-intestinal agents, growth regulators, respiratory stimulants, vitamins, angiotensin converting enzyme inhibitors, anti-asthmatics, anti-cholesterolemics, anti-convulsants, anti-depressants, anti-diarrhea preparations, anti-infective, anti-inflammatory agents, anti-nauseants, anti-stroke agents, anti-tumor drugs, anti-tussives, anti-uricemic drugs, amino-acid preparations, antiemetics, antiobesity drugs, antiparasitics, antipyretics, appetite stimulants, cerebral dilators, chelating agents, cholecystokinin antagonists, cognition activators, deodorants, dermatological agents, diuretics, erythropoietic drugs, fertility agents, synthetic hormones, laxatives, mineral supplements, neuroleptics, neuromuscular agents, peripheral vasodilators, prostaglandins, vaginal preparations, vasoconstrictors, vertigo agents, biguanides, sulphonylurease, meglitinides, PPAR gamma agonist [insulin sensitisers (thiazolidinedione)] and alpha-glucosidase inhibitors.

24. (Original): A dosage form according to claim 21, wherein the high solubility potent active ingredient is selected from the group comprising of benztropine mesylate, distigmine bromide, flupenthixol dihydrochloride, formotexol fumarate, glycopyrrolate, granisetron hydrochloride, bisoprolol fumarate, atropine sulphate, azatadine maleate, carteolol HCl, bromphenaramine maleate, nicotine, oxybutamine chloride, perinodopril erbumine, pilocarpine, poldine methyl sulfate and zalcitane.

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

25. (Original): A dosage form according to claim 1, can be given twice a day or more preferably can be given once a day.

26. (Original): A dosage form according to claim 1, is used for human beings.

27. (Original): A dosage form according to claim 1, wherein the high solubility active ingredient is niacin.

28. (Currently amended): A modified release dosage form according to claim [[1]] 27, wherein the composition of the micromatrix particles and coated micromatrix particles is as follows--Micro matrix particles--Niacin 75% w/w to 99% w/w ~~Eudragit RS~~ EUDRAGIT RS® (Ammonio Methacrylate Copolymer type B USP) 1% w/w to 25% w/w Coated micro matrix particles Micro matrix particles 70% w/w to 99% w/w Hydrogenated castor oil 1% w/w to 30% w/w Magnesium stearate 0% w/w to 2% w/w.

29. (Original): A dosage form according to claim 1, wherein the high solubility active ingredient is sodium valproate.

30. (Original): A process for the preparation of a modified release dosage form comprising a) preparing a micro matrix particles containing high solubility active ingredient and one or more hydrophobic release controlling agent and b) coating the said micromatrix particles containing high solubility active ingredient and one or more hydrophobic release controlling agent.

31. (Original): A modified release dosage form comprising of a metformin hydrochloride prepared by using dual retard technique to control the release of metformin, said dosage

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

form comprising a) micro matrix particles containing metformin hydrochloride and one or more hydrophobic release controlling agent and b) coating of one or more hydrophobic release controlling agent on micro matrix particles.

32. (Original): A dosage form as claimed in claim 31, is in the form of tablet.

33. (Currently amended): A dosage form according to claim 31, wherein the hydrophobic release controlling agents employed for the micro matrix particles are selected from the group comprising ~~consisting of~~ ammonio methacrylate copolymers type A and B as described in USP, methacrylic acid copolymer type A, B and C ~~as described in USP~~, polyacrylate dispersion 30% ~~as described in Ph. Eur.~~, polyvinyl acetate dispersion, ethylcellulose, cellulose acetate, cellulose propionate (lower, medium or higher molecular weight), cellulose acetate propionate, cellulose acetate butyrate, cellulose acetate phthalate, cellulose triacetate, poly(methyl methacrylate), poly(ethyl methacrylate), poly(butyl methacrylate), poly(isobutyl methacrylate), poly(hexyl methacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate), waxes ~~such as~~ selected from the group consisting of beeswax, carnauba wax, microcrystalline wax, and ozokerite; fatty alcohols ~~such as~~ selected from the group consisting of cetostearyl alcohol, stearyl alcohol; cetyl alcohol and myristyl alcohol; and fatty acid esters ~~such as~~ selected from the group consisting of glyceryl monostearate; glycerol monooleate, acetylated monoglycerides, tristearin, tripalmitin, cetyl esters wax, glyceryl palmitostearate, glyceryl behenate and hydrogenated castor oil.

34. (Currently amended): A dosage form according to claim 33, wherein the hydrophobic release controlling agents are selected ~~preferably~~ from ammonio methacrylate copolymers.



Application No: 10/630,348  
Examiner: MERCIER, MELISA S

35. (Currently amended): A dosage form according to claim 34, wherein the ~~preferred from ammonio methacrylate co-polymers preferably from Eudragit RS ammonio methacrylate co-polymers~~ are selected from the groups consisting of ☐ Ammonio Methacrylate Copolymer type B USP[D]], ~~Eudragit RL~~ ☐ Ammonio Methacrylate Copolymer type A USP[D]] and ~~Eudragit NE30D~~ ☐ Polyacrylate dispersion 30% Ph. Eur[☐].

36. (Original): A dosage form according to claim 31, wherein in micro matrix particles, metformin hydrochloride and one or more hydrophobic release controlling agents are present in a ratio of from 100:1 to 100:75.

37. (Currently amended): A dosage form according to claim 36, wherein in micro matrix particles, metformin hydrochloride and one or more hydrophobic release controlling agents are present ~~preferably~~ in a ratio of from 100:2.5 to 100:50.

38. (Currently amended): A dosage form according to claim 36, wherein in micro matrix particles, metformin hydrochloride and one or more hydrophobic release controlling agents are present ~~more preferably~~ in a ratio of from 100:2.5 to 100:30.

39. (Currently amended): A dosage form according to claim 36, wherein in micro matrix particles, metformin hydrochloride and one or more hydrophobic release controlling agents are present ~~most preferably~~ in a ratio of from 100:2.5 to 100:20.

40. (Currently amended): A dosage form according to one of claims 36 ☐ 39, wherein in micro matrix particles, metformin hydrochloride can be less than or equal to 1500 mg.

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

41. (Currently amended): A dosage form according to claim 31, wherein the coating of one or more hydrophobic release controlling agents on said micro matrix particles ~~the hydrophobic release controlling agents employed for coating the said micro matrix~~ are selected from the group ~~comprising~~ consisting ammonio methacrylate copolymers type A and B as described in USP, methacrylic acid copolymer type A, B and C ~~as described in~~ USP, polyacrylate dispersion 30% ~~as described in Ph. Eur.~~ polyvinyl acetate dispersion, ethylcellulose, cellulose acetate, cellulose propionate (lower, medium or higher molecular weight), cellulose acetate propionate, cellulose acetate butyrate, cellulose acetate phthalate, cellulose triacetate, poly(methyl methacrylate), poly(ethyl methacrylate), poly(butyl methacrylate), poly(isobutyl methacrylate), poly(hexyl methacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate), waxes ~~such as selected from the group consisting of~~ beeswax, carnauba wax, microcrystalline wax, and ozokerite; fatty alcohols ~~such as selected from the group consisting of~~ cetostearyl alcohol, stearyl alcohol; cetyl alcohol and myristyl alcohol; and fatty acid esters ~~such as selected from the group consisting of~~ glyceryl monostearate; glycerol monooleate, acetylated monoglycerides, tristearin, tripalmitin, cetyl esters wax, glyceryl palmitostearate, glyceryl behenate and hydrogenated castor oil.

42. (Currently amended): A dosage form according to claim 40, wherein the hydrophobic release controlling agents are ~~is~~ selected from fatty acid esters.

43. (Currently amended): A dosage form according to claim 42, wherein ~~the hydrophobic release controlling agents are~~ fatty acid ester ~~is~~ selected from the group ~~comprising~~ consisting of hydrogenated castor oil and glycerol distearate.

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

44. (Currently amended): A dosage form according to claim 31, wherein micro matrix particles and coating of one or more hydrophobic release controlling agents are present in a ratio of from 100:0.5 to 100:75.

45. (Currently amended): A dosage form according to claim 44, wherein micro matrix particles and coating of one or more hydrophobic release controlling agents are preferably present in a ratio of from 100:1 to 100:50.

46. (Currently amended): A dosage form according to claim 44, wherein micro matrix particles and coating of one or more hydrophobic release controlling agents are more preferably present in a ratio of from 100:2.5 to 100:20.

47. (Currently amended): A modified release dosage form according to claim 31, wherein the composition of the micromatrix particles and coated micromatrix particles is as follows--Micro matrix particles--Metformin hydrochloride 75% w/w to 99% w/w Eudragit RS Ammonio Methacrylate Co-polymer type B USP 1% w/w to 25% w/w Coated micro matrix particles Micro matrix particles 70% w/w to 99% w/w Hydrogenated castor oil 1% w/w to 30% w/w Magnesium stearate 0% w/w to 2% w/w.

48. (Original): A dosage form according to claim 31, wherein the dissolution of metformin hydrochloride is not more than 50% in 1 hour, from 30 to 90% in four hours and not less than 65% in twelve hours.

49. (Original): A dosage form according to claim 31, is once a day oral formulation.

50. (Original): A dosage form according to claim 31, is used for human beings.

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

51. (Original): A dosage form according to claim 31, wherein the maximum plasma metformin concentration is achieved between 700 ng/ml and 2500 ng/ml.
52. (Currently amended): A dosage form according to claim 51, wherein the maximum plasma metformin concentration is achieved preferably between 900 ng/ml and 2400 ng/ml.
53. (Currently amended): A dosage form according to claim 51, wherein the maximum plasma metformin concentration is achieved ~~more preferably~~ between 1000 ng/ml and 2350 ng/ml.
54. (Original): A dosage form according to claim 31, wherein the modified release metformin formulation for once daily administration exhibit invivo mean dissolution time (MDT) of approximately 4 hours to 6 hours.
55. (Original): A dosage form according to claim 31, wherein the minimum plasma metformin concentration (at 24 hours) ranges between 0 and 450 ng/ml after oral administration.
56. (Original): A dosage form as claimed in claim 1, wherein the said dosage form is used to increase the payload of high solubility active ingredient.
57. (Original): A dosage form as claimed in claim 31, wherein the said dosage form is used to increase the payload of metformin hydrochloride.
58. (Original): A dosage form as claimed in claim 1, wherein the said dosage form may optionally contain more than one high solubility active ingredient.

Application No: 10/630,348  
Examiner: MERCIER, MELISA S

59. (Original): A dosage form as claimed in claim 31, wherein the said dosage form may optionally contain more than one antidiabetic active ingredient.

60. (Original): A process for the preparation of a modified release dosage form comprising  
a) preparing a micro matrix particles containing metformin hydrochloride and one or more hydrophobic release controlling agent and b) coating the said micro matrix particles containing metformin hydrochloride and one or more hydrophobic release controlling agent.